

METHODS OF PLANT ROOT EXUDATES ANALYSIS: A REVIEW

P. Dundek, L. Holík, T. Rohlík, L. Hromádko, V. Vranová, K. Rejšek, P. Formánek

Received: January 13, 2011

Abstract

DUNDEK, P., HOLÍK, L., ROHLÍK, T., VRANOVÁ, V., REJŠEK, K., FORMÁNEK, P.: *Methods of plant root exudates analysis: a review*. Acta univ. agric. et silvic. Mendel. Brun., 2011, LIX, No. 3, pp. 241–246

The aim of this review is to summarise current knowledge on methods being used to determine individual compounds and properties of water-soluble plant root exudates. These compounds include amino acids, organic acids and simple sugars, as well as polysaccharides, proteins and organic substances. Qualitative composition of water-soluble root exudates and exudation rate are commonly measured with the aim of consequent synthetic preparation of plant root exudates to be supplied to soil to create artificial rhizosphere for different experimental purposes. Root exudates collection usually requires consequent filtration or centrifugation to remove solids, root detritus and microbial cell debris, and consequent concentration using an evaporator, lyophilizator or ultrafiltration. Methods used for analysis of total groups of compounds (total proteins and total carbohydrates) and total organic carbon are simple. On the other hand, HPLC or GC/MS are commonly used to analyse individual low molecular weight organic molecules (sugars, organic acids and amino acids) with separation using different columns. Other properties such as pH, conductivity or activity of different enzymes as well as gel electrophoresis of proteins are sometimes assessed. All of these methods are discussed in this work.

amino acids, HPLC, GC/MS, organic acids, root exudates, sugars

Substances released by healthy and intact roots into the environment are collectively designated as root exudates. These substances affect the composition of microbial community in immediate proximity of plant roots and processes taking place in rhizosphere. Many compounds are released by plant roots, including inorganic ions and substances, amino acids, amides, sugars, aliphatic acids, aromatic acids, volatile aromatic compounds, gases such as ethylene, vitamins, peptides, proteins, enzymes, plant hormones, alcohols, ketones, olefins, urea, phytoalexins (Vančura, 1988; Grayston *et al.*, 1996; Paynel *et al.*, 2001; Aulakh *et al.*, 2001; Uren, 2007; Neumann and Römheld, 2007). In total, 200 different carbon containing molecules are present in root exudates (Curl and Truelove, 1986), and up to 40% of the net carbon fixed during photosynthesis can be released into the rhizosphere (Whipps and Lynch, 1990). These compounds need to be analysed using different methodological procedures. In this article we present the review of

current methods used for pre-analysis preparation and analysis of different compounds which occur in plant root exudates.

Analysis of plant root exudates

Collection of root exudates

Water-soluble root exudates in collection solutions usually require subsequent concentration on an evaporator or lyophilizator or using ultrafiltration due to low concentration of certain exudate compounds. Not all analysis following exudates collection requires concentrated samples (e. g. total organic carbon or total carbohydrates). Prior to concentration or direct analysis, solids, microorganisms, and root border cells in collected solution should be removed by filtration or centrifugation steps. Depending on the composition of the collection solution, the reduction of sample volume can lead to high salt concentrations,

which may interfere with subsequent analysis, or may even cause irreversible precipitation of certain exudates compounds (e.g. Ca-citrate, Ca-oxalate, proteins). Therefore, if possible, removal of interfering salts using ion-exchange resins prior to sample concentration is recommended (Neumann and Römheld, 2007). Alternatively, solid-phase extraction techniques may be employed for the enrichment of exudates compounds from the diluted trap solution (Johnson *et al.*, 1996; Gerke *et al.*, 1994). High-molecular-weight (HMW) compounds may be concentrated by precipitation with organic solvents (methanol, ethanol, acetone 80% [v/v] for polysaccharides and proteins) or acidification (trichloroacetic acid 10% [w/v], perchloric acid 5% [w/v] for proteins (Neuman *et al.*, 1999). Alternatively, ultrafiltration of the collection solutions or even cultivation of plant roots enclosed in dialysis bags is possible (Sakai and Tadano, 1993).

Analysis of low molecular weight compounds

Before the analysis of concentrated root exudates can be performed separation of exudates into acid, neutral and basic fractions using ion-exchange chromatography followed by separation of organic compounds in each fraction by e.g. HPLC (Schwab *et al.*, 1983; Gransee and Wittenmayer, 2000; Gransee, 2001). Concerning HPLC analysis, Gransee and Wittenmayer (2000) used Bio-Sil NH₂ column (Bio-Rad, München), 150 × 4.6 mm, 5 µm particle size: eluent acetonitrile: water, 85:15 v/v isocratic at 30 °C, flux rate 1.5 mL min⁻¹ for separation of neutral compounds; Schwab *et al.* (1983) used conversion into trimethylsilyl derivatives using Tri-Sil reagent and separated individual carbohydrate derivatives on a GLC fitted with a 200 cm glass column packed with 3% OV-210 on 80- to 100-mesh Chromsorb; Schwab *et al.* (1983) used paper chromatography of underivatized samples of previously separated neutral fraction of root exudates to identify the sugars, using ethylacetate, pyridine and water (8:2:1, v/v/v) as the solvent and 5% *p*-anisidine as the developing agent. An Aminex 87H column, (Bio-Rad, München), 150 × 1.5 mm, 5 µm particle size: eluent 0.005 M sulphuric acid at 30 °C, flux rate 0.6 mL min⁻¹, or a Bio-rad HPX-87 column, with 0.005 M H₂SO₄ following at a rate of 0.5 mL min⁻¹ at ambient temperature as the solvent (UV detection at 210 nm) or mobile phase 0.005 M H₂SO₄, 0.5 mL min⁻¹, 50 °C, wavelength 210 nm were used for separation of organic acids (Schwab *et al.*, 1983; Gransee and Wittenmayer, 2000; Wang *et al.*, 2006); basic compounds were separated after precolumn derivatization with *o*-phthalaldehyde (OPA) reagent on a RP select-B (Merck, Darmstadt), 250 mm × 4 mm, particle size 4 µm, gradient elution with 1 M TBAH in MeOH/MeCN/water 7.5:50: 442.5 → 7.5:350:142.5 (m/m/m), flux rate 1.0 mL min⁻¹ (Gransee and Wittenmayer, 2000). Nardi *et al.* (2002, 2005) used anionic fraction of root exudates to be separated by HPLC on an Aminex column (HPX 87

H – Biorad) with 6 mM H₂SO₄ as mobile phase and detection at 210 nm.

Liu *et al.* (2004) analyzed organic acids using HPLC with an ion-exclusion column (Shim-pack SCR-102h, 0.8 cm × 30 cm). The column was eluted for 30 min. at 23 °C with the mobile phase KH₂PO₄ (18 mM, pH 2.18); the flow rate was 0.8 mL min⁻¹; organic acids were detected at 214 nm and a reversed-phase column C-18 (Silicon 250 mm × 5.6 mm, Shimadzu, Japan) was used. Aulakh *et al.* (2001) analysed organic acids in root exudates according to modified protocol of Badoud and Pratz (1986) when lyophilized exudates were dissolved in distilled water, and derivatized by 1, 4 – dioxan containing O-(4-nitrobenzyl)-N, N-diisopropylisourea. After derivatization at 80 °C for 60 min., the samples were cooled at room temperature and 400 µL H₂O and 600 µL acetonitrile were added. Subsequently, 100 mg Dowex 50 W-X8 (100–200 mesh, p.a.) were added, and the samples were shaken for 2 min., incubated for 15 min at room temperature and centrifuged. The analyse by HPLC was performed using acetonitrile/water gradient (0–20 min, 20–80% acetonitrile) at a flow rate of 1 mL min⁻¹, separation was performed on a reversed-phase octadecyl-silicium column, C-18, 5 µL, 250 × 4.6 mm (Ultrasphere™, Beckman, Germany) with detection at 265 nm. Ma and Miyasaka (1998) searched for oxalate, succinate, malate and citrate in root exudates of taro using HPLC with an anion self-regenerating suppressor (4 mm, model ASRS-I, Dionex). For oxalate and other organic acids analysis Ma and Miyasaka (1998) used an anionic-exchange analytical column and a guard column (both 4 mm, models IonPac AS4A-SC and AG4A-SC, Dionex) with an eluent of 22 mM sodium borate and boric acid at a flow rate of 2.0 mL min⁻¹. The analyse was reconfirmed by the authors using 50 mM NaOH as the eluent for oxalate, succinate, malate, and using 100 mM NaOH for citrate; concentration of organic acids was detected using a conductivity detector. George *et al.* (2002) extracted organic acids from rhizosphere soil by shaking the soil in deionised water with anion exchange strip (recharged with 0.5 M NaHCO₃). After cleaning the soil, elution of organic acids from the strip using 0.5 M HCl, protonation of organic acids by H₂SO₄, consequent methylation with methanol, cooling, application of water, chloroform and shaking, GS/MS analysis of organic acids was used (a Restek Rtx-50 capillary column, 5 m × 0.25 mm i.d., film thickness 0.1 µm, connected to a Hewlett Packard HP 5970 mass spectrum detector, helium as the carrier gas). Vonderheide *et al.* (2006) measured selenium containing amino acids or inorganic compounds using HPLC-ICP-MS. The authors measured selenosulfate SSeO₃⁻² in root exudates using anion-exchange chromatography. Phillips *et al.* (2004) measured amino acids in root solutions by HPLC analysis of fluorescent derivatives produced by reaction with 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate (Cohen and Michaud, 1993). Formánek *et al.* (unpublished)

measured 17 amino acids in root exudates using precolumn derivatization with *o*-phthalaldehyde (OPA). The measurement was performed using an HP 1100 liquid chromatograph (Hewlett Packard, Wilmington, DE, USA) with fluorometric detector FLD HP 1100 operating at 450 nm (Ex = 340 nm) with separation using a Zorbax Exlipse AAA Rapid Resolution (4.6 × 150mm, 3.5 μm particle sizes, Agilent Technologies, USA). A linear gradient profile of mobile phase consisted of 40mM Na₂HPO₄, pH 7.8 (solvent A) and ACN/MeOH/water 45:45:10 (v/v) (solvent B), 0% B (0–1.9 min), 0–57% (1.9–18.1 min), 57–100% (18.1–18.8 min), 100% (18.8–22.3 min), 100–0% (22.3–23.2 min) and 0% (23.2–26 min) was applied at a flow rate of 2.0 mL·min⁻¹ (Formánek *et al.*, 2005). Sometimes, for analysis of amino acids was used Amino Acids analyzer – column elution with sodium citrate buffers (Schwab *et al.*, 1983; Bacilio-Jiménez *et al.*, 2003; Wang *et al.*, 2006). Paynel *et al.* (2001) measured amino acids in root exudates using HPLC as *o*-phthalaldehyde derivatives on a C-18 column using Gold system 8.0. Kravchenko *et al.* (1993; 2004) measured tryptophane in root exudates using HPLC with LiCrosorb RP-18 reversed phase column, which was eluted with solution containing 13.7% acetonitrile and 0.22% acetic acid in water. Wang *et al.* (2006) separated and quantified sugars in root exudates using in HPLC with an RI × 4 detector and Sugar-pak 1. P/N 85188 column (temperature 85 °C, mobile phase Milli-Q water, 0.6 mL·min⁻¹). Bacilio-Jiménez *et al.* (2003) determined carbohydrates in root exudates with the heptafluorobutyrate derivatives of *O*-methylglycosides (after methanolysis in 0.5 M methanol-HCl) by gas chromatography, using a capillary column (25 × 0.32 mm) of 5% silicone OV 210 (gas chromatograph with flame detector, glass solid injector, carrier gas – helium). Erickson *et al.* (2001) used GS/MS for analysis of hydrophobic root exudates, Nardi *et al.* (2005) derivatized root exudates using 2,2-dimethoxypropane *plus* concentrated HCl to obtain corresponding methyl esters; after drying under N flow followed by supplementation of methanol, analysis was done using GS/MS (chromatographic separation using capillary column HP 50, 30 m, film thickness 0.5 μm, internal dia. 0.25).

Other types of analysis

Exudation all of the components released in soil can be suitably evaluated by the amount of carbon (Mench, 1985). Total organic carbon (TOC) is usually determined by wet digestion method (Nelson and Sommers, 1996; Aulakh *et al.*, 2001), or using an

analyzer (Dousset *et al.*, 2001; Personeni *et al.*, 2007; Miya and Firestone, 2001). Sometimes, content of nitrogen (Kjeldahl) is measured (Miya and Firestone, 2001; Nardi *et al.*, 2002). Sometimes dissolved inorganic CO₂ is evacuated from collection media using HCl before the determination of TOC (Personeni *et al.*, 2007). Analysis of total carbohydrates content in root exudates was performed using the anthrone colorimetric assay without lyophilisation of samples (Brink *et al.*, 1960; Aulakh *et al.*, 2001; Formánek *et al.*, unpublished), Miya and Firestone (2001) used phenol-sulphuric acid method to estimate carbohydrate content (Ashwell, 1966; Dubois *et al.*, 1956). Sometimes, exudates are concentrated and carbohydrates in neutral fraction separated from basic and acidic fractions (Schwab *et al.*, 1983). Total soluble sugar content was also determined by colorimetric procedure using sulphonated α -naphthol (Denvor, 1950). For measurement of total sugars Wang *et al.* (2006) hydrolyzed concentrated root exudates with H₂SO₄ under vacuum at 110 °C; after cooling, filtration, drying and dissolution in 0.005 M H₂SO₄, the total sugars were separated and quantified using HPLC. Total amino acids were assessed e.g. by colorimetric procedure using ninhydrin (Spies, 1957); sample of root exudates before analysis by HPLC or amino acid analyzer can also be hydrolysed (Bacilio-Jiménez *et al.*, 2003; Wang *et al.*, 2006) e.g. by 6 M HCl under vacuum at 110 °C and with a drop of phenol to avoid degradation of tyrosine residues as reported by Bacilio-Jiménez *et al.* (2003). Miya and Firestone (2001) determined amino acids in root exudates using a meso-diaminopimelic acid assay (Daniels *et al.*, 1994). Total proteins are mostly determined using Bradford's method (1976) (Bacilio-Jiménez *et al.*, 2003; Nóbrega *et al.*, 2005; Formánek, unpublished). Except frequently measured organic compounds, pH of the rhizosphere or conductivity of root exudates is sometimes measured. Wang *et al.* (2006) measured conductivity of root exudates at room temperature and after being exposed to high temperature, and from these values they calculated the percentage of electrolytes osmosized from roots. Activity of different enzymes (haem-containing peroxidases, unspecific copper-containing oxidases, unspecific haem-containing oxidase, flavin-containing monooxygenases – hydroxylating, aromatic-ring cleavage dioxygenases, oxydases that catalyse H₂O₂ formation and bleaching of dye macromolecules) or gel electrophoresis of proteins is sometimes assessed (Gramss and Rudeschko, 1998).

CONCLUSION

Determination of three most abundant compounds in water-soluble plant root exudates (sugars, organic acids and amino acids) is highly needed for various purposes. These compounds (even dominant within root exudates) occur in very low concentrations and sensitive analytical procedures

are necessary for their determination. Different approaches with the use of HPLC or GS/MS are utilized to determine individual sugars, organic acids and amino acids in plant root exudates. Determination of other properties of plant root exudates (concentration of total carbohydrates, proteins, pH, conductivity or activity of different enzymes as well as gel electrophoresis of proteins) is sometimes used.

The study was supported by the grant MSM6215648902 / Forest and Wood: the support of functionally integrated forest management and use of wood as a renewable raw material phase 4/2/3, part II "The management strategy of nature conservation areas", and by project "Extension of current knowledge on bioavailable amino acids in soil and their utilization by soil microorganisms and plant roots" (IGA FFWT MENDELU No. 47/2010–2012).

REFERENCES

- ASHWELL, G., 1966: New colorimetric methods of sugar analysis. In *Methods in enzymology*, Vol. VIII, E. F. Neufeld, V. Ginsburg (Eds). pp 85–95. Academic Press, New York. ISBN 978-0-12-181808-1.
- AULAKH, M. S., WASSMANN, R., BUENO, C., KREUZWIESER, J., RENNENBERG, H., 2001: Characterization of root exudates at different growth stages of ten rice (*Oryza sativa* L.) cultivars. *Plant Biology* 3, 2: 139–148. ISSN 1438-8677.
- BACILIO-JIMENEZ, M., AGUILAR-FLORES, S., VENTURA-ZAPATA, E., PÉREZ-CAMPOS, E., BOUQUELET, S., ZENTENO, E., 2003: Chemical characterization of root exudates from rice (*Oryza sativa*) and their effect on the chemotactic response of endophytic bacteria. *Plant and Soil* 249, 2: 271–277. ISSN 0032-079X.
- BADOUD, R., PRATZ, G., 1986: Improved high-performance liquid chromatographic analysis of some carboxylic acids in food and beverages as their p-nitrobenzyl esters. *Journal of Chromatography* 360, 119–136. ISSN 0021-9673.
- BRADFORD, M. M., 1976: A rapid and sensitive method for the quantitation of microgram quantities of protein utilising the principle of protein-dye binding. *Analytical Biochemistry* 72, 2: 248–254. ISSN 0003-2697.
- BRINK, R. H. J., DUBACH, P., LYNCH, D. L., 1960: Measurement of carbohydrates in soil hydrolyzates with anthrone. *Soil Science* 89, 3: 157–166. ISSN 0038-075X.
- COHEN, S. A., MICHAUD, D. P., 1993: Synthesis of a fluorescent derivatizing reagent, 6-aminoquinolyl-N-hydroxysuccinimidyl carbamate, and its application for the analysis of hydrolysate amino acids via high-performance liquid chromatography. *Analytical Biochemistry* 211, 2: 279–287. ISSN 0003-2697.
- CURL, E. A., TRUELOVE, B., 1986: *The Rhizosphere*. Springer-Verlag, New York. Springer-Verlag, Berlin-Heidelberg-New York-Tokyo, 288 p. ISBN 3-540-15803-0.
- DANIELS, L., HANSON, R. S., PHILIPS, J. A., 1994: Chemical analysis. In: Gerhard, P. et al. (Eds). *Method for general and molecular bacteriology*. American Society for Microbiology, Washington, DC, 512–554.
- DENVOR, A. W., 1950: Carbohydrate tests using sulphonated α -naphthol. *Journal of the American Chemical Society* 72, 2008–2012.
- DOUSSET, S., MOREL, J. L., JACOBSON, A., BITTON, G., 2001: Copper binding capacity of root exudates of cultivated plants and associated weeds. *Biology and Fertility of Soils* 34, 4: 230–234. ISSN 0178-2762.
- DUBOIS, M., GILLES, K. A., HAMILTON, J. K., REBERS, P. A., SMITH, F., 1956: Colorimetric method for determination of sugars and related substances. *Analytical Chemistry* 28, 3: 350–356. ISSN 0003-2700.
- ERICKSON, J., SCHOTT, D., REVERRI, T., MUHSIN, W., RUTTLEDGE, T., 2001: GC-MS analysis of hydrophobic root exudates of *Sorghum* and implications on the parasitic plant *Striga asiatica*. *Journal of Agricultural and Food Chemistry* 49, 5537–5542. ISSN 0021-8561.
- FORMÁNEK, P., KLEJDUS, B., VRANOVÁ, V., 2005: Bio-available amino acids extraction from soil by demineralized water and 0.5M ammonium acetate. *Amino Acids* 28, 4: 427–429. ISSN 0939-4451.
- GEORGE, T. S., GREGORY, P. J., WOOD, M., READ, D., BURESH, R. J., 2002: Phosphatase activity and organic acids in the rhizosphere of potential agroforestry species and maize. *Soil Biology and Biochemistry* 34, 10: 1487–1494. ISSN 0038-0717.
- GERKE, J., RÖMER, W., JUNK, A., 1994: The excretion of citric and malic acid by proteoid roots of *Lupinus albus* L.: effect on soil solution concentrations of phosphate, iron and aluminium in the proteoid rhizosphere sample of an oxisol and a luvisol. *Journal of Plant Nutrition and Soil Science* 157, 4: 289–294. ISSN 1436-8730.
- GRAMSS, G., RIDESCHKO, O., 1998: Activities of oxidoreductase enzymes in tissue extracts and sterile root exudates of three crop plants, and some properties of the peroxidase component. *New Phytologists* 138, 3: 401–409. ISSN 0028-646X.
- GRANSEE, A., 2001: Effects of root exudates on nutrient availability in the rhizosphere. In:

- Horst, W. J. et al. 2001. *Plant nutrition-food security and sustainability of agro-ecosystems*, 626–627. Food security and sustainability and sustainability of agro-ecosystems through basic and applied research. ISBN 0792371054.
- GRANSEE, A., WITTENMAYER, L., 2000: Qualitative and quantitative analysis of water-soluble root exudates in relation to plant species and development. *Journal of Plant Nutrition and Soil Science* 163, 4: 381–385. ISSN 1436-8730.
- GRAYSTON, S. J., VAUGHAN, D., JONES, D., 1996: Rhizosphere carbon flow in trees, in comparison with annual plants: the importance of root exudation and its impact on microbial activity and nutrient availability. *Applied Soil Ecology* 5, 1: 29–56. ISSN 0929-1393.
- JOHNSON, J. F., ALLAN, D. L., VANCE, C. P., WEIBLEN, G., 1996: Root Carbon Dioxide Fixation by Phosphorus-Deficient *Lupinus albus* (Contribution to Organic Acid Exudation by Proteoid Roots). *Plant Physiology* 112, 1: 19–30. ISSN 0032-0889.
- KRAVCHENKO, L. V., AZAROVA, T. S., MAKAROVA, N. M., TIKHONOVICH, I. A., 2004: The effect of tryptophan present in plant roots exudates on the phyto-stimulating activity of rhizobacteria. *Microbiology* 73, 2: 156–158. ISSN 0026-2617.
- KRAVCHENKO, L. V., LEONOVA, E. I., 1993: The use of the root exometabolite tryptophan by plant-associated bacteria for the biosynthesis of indole-3-acetic acid. *Microbiologiya* 62, 3: 453–459. ISSN 0026-3656.
- LI, Y. G., LI, L. H., JIANG, G. M., NIU, S. L., LIU, M. Z., GAO, L. M., PENG, Y., JIANG, C. D., 2004: Traits of chlorophyll fluorescence in 99 plant species from the sparse-elm grassland in Hunshandak Sandland. *Photosynthetica* 42, 2: 243–249. ISSN 0300-3604.
- LIN, C. H., LERCH, R. N., GARRETT, H. E., GEORGE, M. F., 2004: Incorporating forage grasses in riparian buffers for bioremediation of atrazine, isoxaflutole and nitrate in Missouri. *Agroforestry Systems* 63, 1: 91–99. ISSN 0167-4366.
- MA, Z., MIYASAKA, S. C., 1998: Oxalate exudation by taro in response to Al. *Plant Physiology* 118, 5: 861–865. ISSN 0032-0889.
- MENCH, M., 1985: Influence des exudats racinaires solubles sur la dynamique des métaux dans la rhizosphère du maïs (*Zea mays* L.). Thèse de doctorat, University of Nancy. FRBNF 37594962.
- MIYA, R. K., FIRESTONE, M. K., 2001: Enhanced phenanthrene biodegradation in soil by slender oat root exudates and root debris. *Journal of Environmental Quality* 30, 6: 1911–1918. ISSN 0047-2425.
- NARDI, S., SESSI, E., PIZZEGHELLO, D., STURARO, A., RELLA, R., PARVOLI, G., 2002: Biological activity of soil organic matter mobilized by root exudates. *Chemosphere* 46, 7: 1075–1081. ISSN 0045-6535.
- NARDI, S., TOSONI, M., PIZZEGHELLO, D., PROVENZANO, M. R., CILENTI, A., STURARO, A., RELLA, R., VIANELLO, A., 2005: Chemical characteristics and biological activity of organic substances extracted from soils by root exudates. *Soil Science Society of America Journal* 69, 2012–2019. ISSN 0361-5995.
- NELSON, D. W., SOMMERS, L. E., 1996: Total carbon, organic carbon, and organic matter. In: Sparks, D. L. (Ed.), *Methods of Soil Analysis, Chemical Methods*, Vol. 5, part 3. Soil Science Society of America Inc., Madison, WI, USA, 961–1010. ISSN 0361-5995.
- NEUMAN, G., HÜLSTER, A., RÖMHELD, V., 1999: PCDD/PCDF-mobilizing compounds in root exudates of zucchini. *Organohalogen Compounds* 41: 331–334. ISSN 1026-4892.
- NEUMAN, G., RÖMHELD, V., 2007: The release of root exudates as affected by the plant physiological status. In: Pinton, R., Varanini, Z., Nannipieri, P. (Eds.), *The rhizosphere: Biochemistry and organic substances at the soil-plant interface*, 23–72. ISBN 9780849338557.
- NÓBREGA, F. M., SANTOS, I. S., DA DUNHA, M., CARVALHO, A. O., GOMES, V. M., 2005: Antimicrobial proteins from cowpea root exudates: inhibitory activity against *Fusarium oxysporum* and purification of chitinase-like proteins. *Plant and Soil* 272, 1–2: 223–232. ISSN 0032-079X.
- PAYNEL, F., MURRAY, J., CLIQUET, J. B., 2001: Root exudates: a pathway for short-term N transfer from clover and ryegrass. *Plant and Soil* 229, 2: 235–243. ISSN 0032-079X.
- PERSONENI, E., NGUYEN, C., MARCHAL, P., PAGÉS, L., 2007: Experimental evaluation of an efflux-influx model of C exudation by individual apical root segments. *Journal of Experimental Botany* 58, 8: 2091–2099. ISSN 0022-0957.
- PHILIPS, D. A., FOX, T. C., KING, M. D., BHUVANESWARI, T. V., TEUBER, L. R., 2004: Microbial products trigger amino acids exudation from plant roots. *Plant Physiology* 136, 1: 2887–2894. ISSN 0032-0889.
- SAKAI, H., TADANO, T., 1993: Characteristics of response of acid phosphatase secreted by the roots of several crops to various conditions in the grow media. *Soil Science and Plant Nutrition* 39, 3: 437–444. ISSN 0038-0768.
- SCHWAB, S. M., MENGE, J. A., LEONARD, R. T., 1983: Quantitative and qualitative effects of phosphorus on extracts and exudates of sudangrass roots in relation to vesicular-arbuscular mycorrhiza formation. *Plant Physiology* 73, 3: 761–765. ISSN 0032-0889.
- SPIES, J. R., 1957: Colorimetric procedures for amino acids. In: Colowick, S. P., Kaplan, N. O. *Methods in Enzymology*, Vol. III. pp. 468–471. Academic Press, New York.
- UREN, N. C., 2007: Type, amounts, and possible functions of compounds released into the rhizosphere by soil-grown plants. In: Pinton, R., Varanini, Z., Nannipieri, P. (Eds.), *The rhizosphere: Biochemistry and organic substances at the soil-plant interface*, pp 1–21, CRC Press, Boca Raton, FL.

- VANČURA, V., 1988: Plant metabolites in soil. In: Vančura, V., Kunc, F. (Eds.). *Soil microbial associations: control of structures and functions*. Academia Praha, 57–144.
- VONDERHEIDE, A. P., MOUNICOU, S., MEIJA, J., HENRY, H. F., CARUSO, J. A., SHANN, J. R., 2006: Investigation of selenium-containing root exudates of *Brassica juncea* using HPLC-ICP-MS and ESI-qTOF-MS. *Analyst* 131, 33–40. ISSN 0003-2654.
- WANG, P., BI, S., WANG, S., DING, Q., 2006: Variation of wheat root exudates under aluminium stress. *Journal of Agricultural and Food Chemistry* 54, 10040–10046. ISSN 0021-8561.
- WHIPPS, J. M., LYNCH, J. M., 1990: Substrate flow in the rhizosphere. *Plant and Soil* 129, 1: 1–10. ISSN 0032-079X.

Address

Ing. Peter Dundek, Bc. Ladislav Holík, Bc. Tomáš Rohlík, Ing. Ladislav Hromádko, Ing. Valerie Vranová, Ph.D., doc. Ing. Klement Rejšek, CSc., doc. RNDr. Pavel Formánek, Ph.D., Ústav geologie a pedologie, Mendelova univerzita v Brně, Zemědělská 3, 613 00 Brno, Česká republika, e-mail: xdundek@node.mendelu.cz, ladisholik@seznam.cz, toro83@centrum.cz, xhromad4@node.mendelu.cz, vranova@mendelu.cz, kr@mendelu.cz., formanek@mendelu.cz